

[https://data.fulir.irb.hr/hr/islandora/search?display=default&f%5B0%5D=%2DRELS\\_EXT\\_hasModel\\_uri\\_s%3A%22info%3Afedora/islandora%3AcollectionCModel%22&f%5B1%5D=mods\\_language\\_ms%3A%22eng%22&f%5B2%5D=mods\\_keywords\\_ms%3A%22MeV%20TOF%20SIMS%22&sort=score%20desc&islandora\\_solr\\_search\\_navigation=0](https://data.fulir.irb.hr/hr/islandora/search?display=default&f%5B0%5D=%2DRELS_EXT_hasModel_uri_s%3A%22info%3Afedora/islandora%3AcollectionCModel%22&f%5B1%5D=mods_language_ms%3A%22eng%22&f%5B2%5D=mods_keywords_ms%3A%22MeV%20TOF%20SIMS%22&sort=score%20desc&islandora_solr_search_navigation=0)

Vrijeme izvoza: 19.04.2024. 11:05:26

Repozitorij: data.fulir.irb.hr

Ukupan broj zapisa na URL-u: 2

Broj izvezenih zapisa: 2

| Naslov                                                                                                          | URL                                                                                             | Autori                                                                         | Naslov izvornika |
|-----------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------|------------------|
| MeV TOF SIMS determination of deposition order between optically distinguishable and indistinguishable inks     | <a href="https://urn.nsk.hr/urn:nbn:hr:241:440276">https://urn.nsk.hr/urn:nbn:hr:241:440276</a> | Barac, Marko; Bogdanović-Radović, Ivančica; Brajković, Marko; Siketić, Zdravko |                  |
| Dependence of MeV TOF SIMS secondary molecular ion yield from phthalocyanine blue on primary ion stopping power | <a href="https://urn.nsk.hr/urn:nbn:hr:241:471554">https://urn.nsk.hr/urn:nbn:hr:241:471554</a> | Brajković, Marko; Siketić, Zdravko; Bogdanović-Radović, Ivančica; Barac, Marko |                  |