

[https://data.fulir.irb.hr/hr/islandora/search?display=default&f%5B0%5D=%2DRELS\\_EXT\\_hasModel\\_uri\\_s%3A%22info%3Afedora/islandora%3AcollectionCModel%22&f%5B2%5D=mods\\_keywords\\_ms%3A%22MeV%20TOF%20SIMS%22&sort=score%20desc&islandora\\_solr\\_search\\_navigation=0](https://data.fulir.irb.hr/hr/islandora/search?display=default&f%5B0%5D=%2DRELS_EXT_hasModel_uri_s%3A%22info%3Afedora/islandora%3AcollectionCModel%22&f%5B2%5D=mods_keywords_ms%3A%22MeV%20TOF%20SIMS%22&sort=score%20desc&islandora_solr_search_navigation=0)

Vrijeme izvoza: 10.08.2022. 21:56:45

Repozitorij: data.fulir.irb.hr

Ukupan broj zapisa na URL-u: 2

Broj izvezenih zapisa: 2

<b>Naslov</b>	<b>URL</b>	<b>Autori</b>	<b>Naslov izvornika</b>
Dependence of MeV TOF SIMS secondary molecular ion yield from phthalocyanine blue on primary ion stopping power	<a href="https://urn.nsk.hr/urn:nbn:hr:241:471554">https://urn.nsk.hr/urn:nbn:hr:241:471554</a>	Brajković, Marko; Siketić, Zdravko; Bogdanović-Radović, Ivančica; Barac, Marko	
MeV TOF SIMS determination of deposition order between optically distinguishable and indistinguishable inks	<a href="https://urn.nsk.hr/urn:nbn:hr:241:440276">https://urn.nsk.hr/urn:nbn:hr:241:440276</a>	Barac, Marko; Bogdanović-Radović, Ivančica; Brajković, Marko; Siketić, Zdravko	